

Seat No.: _____

Enrolment No. _____

GUJARAT TECHNOLOGICAL UNIVERSITY
BE SEMESTER 1st / 2nd (OLD) EXAMINATION WINTER 2016

Subject Code: 110009

Date: 20/01/2017

Subject Name: Mathematics-II

Time: 10:30 AM TO 1:30 PM

Total Marks: 70

Instructions:

1. Attempt any five questions.
2. Make suitable assumptions wherever necessary.
3. Figures to the right indicate full marks.

- Q.1 (a) 1. Solve the following system by Gauss elimination method: 03
- $$2x_1 + 2x_2 + 2x_3 = 0, -2x_1 + 5x_2 + 2x_3 = 1, 8x_1 + x_2 + 4x_3 = -1$$
2. Solve the system 04
- $$2x_1 - x_2 = \lambda x_1, 2x_1 - x_2 + x_3 = \lambda x_2, -2x_1 + 2x_2 + x_3 = \lambda x_3$$
- for x_1, x_2, x_3 in the two cases $\lambda = 1$ and $\lambda = 2$.
- (b) 1. Solve the system 04
- $$-2y + 3z = 1; 3x + 6y - 3z = -2; 6x + 6y + 3z = 5$$
- by Gauss-Jordan method.
2. Find the rank of the matrix $\begin{bmatrix} 1 & -1 & 3 & 6 \\ 1 & 3 & -3 & -4 \\ 5 & 3 & 3 & 11 \end{bmatrix}$. 02
3. Prove that for the matrix $A = \begin{bmatrix} -1 & 2+i & 5-3i \\ 2-i & 7 & 5i \\ 5+3i & -5i & 2 \end{bmatrix}$, iA is a skew Hermitian matrix. 01
- Q.2 (a) 1. Find the vectors in R^2 with Euclidean norm 1 whose Euclidean inner product with $(3, -1)$ is zero. 03
2. State Cauchy-Schwarz inequality in R^n . Verify Cauchy-Schwarz inequality for the vectors $u = (0, -2, 2, 1)$ and $v = (-1, -1, 1, 1)$. Also, find $\|u - v\|$. 04
- (b) 1. Let $T: R^3 \rightarrow R^3$ be a multiplication by A . Determine whether T has an inverse. If so, find $T^{-1} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$ where $A = \begin{bmatrix} 1 & 4 & -1 \\ 1 & 2 & 1 \\ -1 & 1 & 0 \end{bmatrix}$ 02

2. Show that the linear transformation $T: R^2 \rightarrow R^2$ defined by $T(x, y) = (2x - 6y, -5x + 15y)$ is not invertible. 02
3. Determine the algebraic and geometric multiplicity of the matrix $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & -3 & 3 \end{bmatrix}$ 03
- Q.3 (a) 1. Let R^3 have the Euclidean inner product. Transform the basis $u_1 = (1, 1, 1), u_2 = (1, -2, 1), u_3 = (1, 2, 3)$ into an orthogonal basis using the Gram-Schmidt process. 03
2. Find the least square solution of the linear system $AX = b$ given by $x_1 - x_2 = 4, 3x_1 + 2x_2 = 1, -2x_1 + 4x_2 = 3$ and find orthogonal projection of b on the column space of A . 04
- (b) 1. Verify Cayley-Hamilton theorem for the matrix $A = \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$ and hence find A^{-1} and A^4 . 03
2. Reduce the quadratic form $Q(x, y, z) = 3x^2 + 3z^2 + 4xy + 8yz + 8xz$ to canonical form by linear transformation. 04
- Q.4 (a) 1. Show that $S = \{1 - x - x^3, -2 + 3x + x^2 + 2x^3, 1 + x^2 + 5x^3\}$ is linearly independent in P_3 . 03
2. Find the inverse of the matrix $A = \begin{bmatrix} 2 & 1 & 1 \\ 3 & 2 & 1 \\ 2 & 1 & 2 \end{bmatrix}$ by Gauss-Jordan elimination method. 04
- (b) 1. Show that $S = \left\{ \begin{bmatrix} 1 & 2 \\ 1 & -2 \end{bmatrix}, \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 2 \\ 3 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ -1 & 2 \end{bmatrix} \right\}$ is a basis for M_{22} . 03
2. Find the matrix representation of quadratic form $x^2 - 8y^2 + 14z^2 - 6xy + 10yz + 4xz$ 02
3. Obtain the matrix of a linear transformation $T: R^2 \rightarrow R^3$ defined by the formula $T(x_1, x_2) = (x_1 + x_2, 2x_1 - x_2, 7x_2)$ with respect to the standard basis $B_1 = \{u_1, u_2\}$ and $B_2 = \{v_1, v_2, v_3\}$. 02
- Q.5 (a) 1. Find the eigenvalues and eigenvectors of the matrix $A = \begin{bmatrix} 3 & 1 & 4 \\ 0 & 2 & 6 \\ 0 & 0 & 5 \end{bmatrix}$. 04

2. Find matrix P that diagonalizes the matrix $A = \begin{bmatrix} 1 & 0 \\ 6 & -1 \end{bmatrix}$. Also determine $P^{-1}AP$. 03

- (b) 1. Let $T: R^2 \rightarrow R^3$ the linear transformation defined by 04

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_2 \\ -5x_1 + 13x_2 \\ -7x_1 + 16x_2 \end{bmatrix}. \text{ Find the matrix for the transformation}$$

T with respect to the basis $B_1 = \{u_1, u_2\}$ for R^2 and $B_2 = \{v_1, v_2, v_3\}$ for

$$R^3, \text{ where } u_1 = \begin{bmatrix} 3 \\ 1 \end{bmatrix}, u_2 = \begin{bmatrix} 5 \\ 2 \end{bmatrix}, v_1 = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, v_2 = \begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix}, v_3 = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix}$$

2. Consider the basis $S = \{v_1, v_2, v_3\}$ for R^3 where $v_1 = (1, 1, 1)$, $v_2 = (1, 1, 0)$ and $v_3 = (1, 0, 0)$ and let $T: R^3 \rightarrow R^3$ be the linear operator such that $T(v_1) = (2, 1, -4)$, $T(v_2) = (3, 0, 1)$, $T(v_3) = (-1, 5, 1)$. Find a formula for $T(x_1, x_2, x_3)$ and use the formula to find $T(2, 4, -1)$. 03

- Q.6 (a) 1. Let V be the set of all ordered pairs of real numbers with vector addition defined as 04

$(x_1 + y_1) + (x_2 + y_2) = (x_1 + x_2 + 1, y_1 + y_2 + 1)$ show that first five axioms for vector addition are satisfied. Clearly mention the zero vector and additive inverse.

2. State Dimension Theorem and verify that for the given matrix A 03

$$A = \begin{bmatrix} 2 & 2 & -1 & 0 & 1 \\ -1 & -1 & 2 & -3 & 1 \\ 1 & 1 & -2 & 0 & -1 \\ 0 & 0 & 1 & 1 & 1 \end{bmatrix}$$

- (b) 1. Let $S = \{(2, 0, 0), (0, 2, 0)\}$ show that S is not a basis for R^3 even if it is linearly independent. 03

2. If $V = \{(x, y, z) / x - 3y + 4z = 0, x, y, z \in R\}$, prove that V is a subspace of a vector space of R^3 . 04

- Q.7 (a) 1. Consider a basis $S = \{b_1, b_2\}$ for R^2 , where $b_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $b_2 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$. 03

Suppose v in R^2 has the coordinate vector $[v]_S = \begin{bmatrix} -2 \\ 3 \end{bmatrix}$. Find v .

2. Let V be the space spanned by $v_1 = \sin x$, $v_2 = \cos x$, $v_3 = x$. Show that $S = v_1, v_2, v_3$ forms a basis for V . 01

3 Find a basis for column-space of the matrix $A = \begin{bmatrix} 1 & 2 & -1 & 3 \\ -2 & -4 & 4 & -7 \\ 1 & 2 & 1 & 2 \end{bmatrix}$. 03

(b) 1. Let $T: R^2 \rightarrow R^2$ be the linear operator defined by 04

$$T\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1 - 2x_2 \\ -x_2 \end{bmatrix} \text{ and } s_1 = \{v_1, v_2\} \text{ and } s_2 = \{w_1, w_2\} \text{ where}$$

$$v_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, v_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, w_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, w_2 = \begin{bmatrix} -3 \\ 4 \end{bmatrix}.$$

(i) Find the matrix T w.r.t. the basis s_1

(ii) Find the matrix T w.r.t. the basis s_2

2. Define: Symmetric matrix, Skew- Symmetric Matrix, Diagonal matrix 03
