

**GUJARAT TECHNOLOGICAL UNIVERSITY**  
**BE SEMESTER 1<sup>st</sup> / 2<sup>nd</sup> (NEW) EXAMINATION WINTER 2016**

**Subject Code: 2110015****Date: 20/01/2017****Subject Name: Vector Calculus and Linear Algebra****Time: 10:30 AM TO 1:30 PM****Total Marks: 70****Instructions:**

1. Question No. 1 is compulsory. Attempt any four out of remaining Six questions.
2. Make suitable assumptions wherever necessary.
3. Figures to the right indicate full marks.

- | <b>Q.1</b> | <b>Objective Question (MCQ)</b>  | <b>Mark</b> |
|------------|--|-------------|
|            | <b>(a) Choose the appropriate answer for the following MCQs.</b>   | <b>07</b>   |
| 1.         | Rank of $3 \times 3$ invertible matrix is  |             |
|            | a) 1                      b) 2                      c) 3                      d) 4                                       |             |
| 2.         | Let $A$ be a Skew- Hermitian matrix then $A =$ _____   |             |
|            | a) $A^T$ b) $(\overline{A})^T$ c) $-A^T$ d) $-(\overline{A})^T$  |             |
| 3.         | The set $S = \{1, x, x^2, x^3\}$ span which of the following?  |             |
|            | a) $P_3$ b) $R$ c) $R^3$ d) $M_{33}$   |             |
| 4.         | Let $u = (1, -2)$ be a vector in $R^2$ with the Euclidean inner product, then $\ u\ $ is                                 |             |
|            | a) 1                      b) 5                      c) $\sqrt{5}$ d) $\sqrt{3}$  |             |
| 5.         | If $A$ is a matrix with 6 columns and $rank(A) = 2$ , then $Nullity(A)$ is   |             |
|            | a) 2                      b) 4                      c) 0                      d) None of these                           |             |
| 6.         | If $\lambda_1 = 2, \lambda_2 = 6$ are the eigen values of the matrix $A$ , then the eigen values of $A^T$ are            |             |
|            | a) 2 & 6                      b) $\frac{1}{2}$ & $\frac{1}{6}$ c) 4 & 36                      e) None of these           |             |
| 7.         | The product of the eigen values of matrix $A = \begin{bmatrix} 4 & 2 & -2 \\ -5 & 3 & 2 \\ -2 & 4 & 1 \end{bmatrix}$ is, |             |

**(b) Choose the appropriate answer for the following MCQs.**

1. Let  $I : V \rightarrow V$  be an identity operator, then  $\ker(I)$  is,
 

a)  $V$                       b)  $\{0\}$                       c) 0                      d) None of these
2. The mapping  $T : R^3 \rightarrow R^3$  defined by  $T(x, y, z) = (x, y, 0)$  is called as
 

a) Projection                      b) Reflection  
c) Rotation                      d) Magnification
3. If  $f_1 = x$  and  $f_2 = \sin x$ , then Wronskian  $w\left(\frac{\pi}{2}\right) = \underline{\hspace{2cm}}$ 

a)  $\frac{\pi}{2}$                       b) 1                      c) 0                      d) -1
4. If  $\vec{r} = xi + yj + zk$ , then divergence of  $\vec{r}$  is
 

a) 2                      b) -2                      c) 3                      d) -3
5. The value of  $\text{curl}(\text{grad } \phi)$ , where  $\phi = 2x^2 - 3y^2 + 4z^2$  is
 

a)  $4xi - 6yj + 8zk$                       b)  $4x - 6y + 8z$                       c) 6                      d)  $\vec{0}$
6. The weighted Euclidean inner product  $\langle u, v \rangle = 3u_1v_1 + 2u_2v_2$  is the inner product on  $R^2$  generated by
 

a)  $\begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix}$                       b)  $\begin{bmatrix} \sqrt{3} & 0 \\ 0 & \sqrt{2} \end{bmatrix}$                       c)  $\begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$                       d)  $\begin{bmatrix} 0 & 2 \\ 3 & 0 \end{bmatrix}$
7. If  $V$  is a finite-dimensional vector space, and  $T : V \rightarrow V$  is a linear operator and  $\ker(T) = \{0\}$ , then
 

a)  $R(T) \neq V$                       b)  $T$  is one-to-one  
c)  $\text{Nullity}(T) \neq 0$                       d) None of these

**Q.2 (a) Convert the following matrix in to reduced row echelon form and hence find the rank of a matrix. 03**

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 4 & 2 \\ 2 & 6 & 5 \end{bmatrix}$$

**(b) Solve the following system of equations by Gauss Elimination method 04**

$$\begin{aligned} x_1 - 2x_2 + 3x_3 &= -2 \\ -x_1 + x_2 - 2x_3 &= 3 \\ 2x_1 - x_2 + 3x_3 &= -7 \end{aligned}$$

- (c) (i) Find the inverse of the matrix  $A = \begin{bmatrix} 1 & 0 & 1 \\ -1 & 1 & 1 \\ 0 & 1 & 0 \end{bmatrix}$  using Gauss Jordan 03

Method

- (ii) For what choices of parameter the following system is consistent. 04

$$x_1 + x_2 + 2x_3 + x_4 = 1$$

$$x_1 + 2x_3 = 0$$

$$2x_1 + 2x_2 + 3x_3 = \lambda$$

$$x_2 + x_3 + 3x_4 = 2\lambda$$

- Q.3** (a) Consider the vectors  $u = (1, 2, -1)$  and  $v = (6, 4, 2)$  in  $R^3$ . Show that  $w = (9, 2, 7)$  is a linear combination of  $u$  and  $v$ . 03

- (b) Determine a basis for and the dimension of the solution space of the homogeneous system 04

$$2x_1 + 2x_2 - x_3 + x_5 = 0$$

$$-x_1 - x_2 + 2x_3 - 3x_4 + x_5 = 0$$

$$x_1 + x_2 - 2x_3 - x_5 = 0$$

$$x_3 + x_4 + x_5 = 0$$

- (c) Show that the set of all pairs of real numbers of the form  $(1, x)$  with the operations  $(1, y) + (1, y') = (1, y + y')$  and  $k(1, y) = (1, ky)$  is a vector space. 07

- Q.4** (a) Sketch the unit circle in an  $xy$ -coordinate system in  $R^2$  using 03

1. The Euclidean inner product  $\langle u, v \rangle = u_1v_1 + u_2v_2$ .

2. The weighted Euclidean inner product  $\langle u, v \rangle = \frac{1}{9}u_1v_1 + \frac{1}{4}u_2v_2$ .

- (b) Attempt the following. 04

1. Let  $u = (u_1, u_2)$  and  $v = (v_1, v_2)$  be vectors in  $R^2$ . Verify that the weighted Euclidean inner product  $\langle u, v \rangle = 3u_1v_1 + 2u_2v_2$  satisfies the four inner product axioms.

2. Let  $R^4$  have the Euclidean inner product. Find the cosine of the angle  $\theta$  between the vectors  $u = (4, 3, 1, -2)$  and  $v = (-2, 1, 2, 3)$ .

- (c) Consider the vector space  $R^3$  with the Euclidean inner product. Apply the Gram-Schmidt process to transform the basis vectors  $u_1 = (1, 1, 1)$ ,  $u_2 = (0, 1, 1)$  &  $u_3 = (0, 0, 1)$  into an orthogonal basis  $\{v_1, v_2, v_3\}$ ; then normalize the orthogonal basis vectors to obtain an orthonormal basis  $\{q_1, q_2, q_3\}$ . 07

- Q.5** (a) Let  $T : P_1 \rightarrow P_2$  be the linear transformation defined by  $T(p(x)) = x(p(x))$ . 03

Find the matrix for  $T$  with respect to the standard bases  $B = \{u_1, u_2\}$  and  $B' = \{v_1, v_2, v_3\}$ , where  $u_1 = 1$ ,  $u_2 = x$ ;  $v_1 = 1$ ,  $v_2 = x$ ,  $v_3 = x^2$ .

- (b) Express  $\begin{bmatrix} 4 + 2i & 7 & 3 - i \\ 0 & 3i & -2 \\ 5 + 3i & -7 + i & 9 + 6i \end{bmatrix}$  as the sum of a Hermitian and a skew-Hermitian matrix. 04

- (c) State the Dimension theorem for Linear Transformation and find the rank and nullity 07

of  $T_A$ , where  $T_A : R^6 \rightarrow R^4$  be multiplication by

$$A = \begin{bmatrix} -1 & 2 & 0 & 4 & 5 & -3 \\ 3 & -7 & 2 & 0 & 1 & 4 \\ 2 & -5 & 2 & 4 & 6 & 1 \\ 4 & -9 & 2 & -4 & -4 & 7 \end{bmatrix}$$

**Q.6 (a)** Determine the algebraic and geometric multiplicity of  $\begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$ . **03**

**(b)** For the matrix  $A = \begin{bmatrix} -4 & -6 \\ 3 & 5 \end{bmatrix}$ , find the nonsingular matrix  $P$  and the diagonal matrix  $D$  such that  $D = P^{-1}AP$ . **04**

**(c)** Determine  $A^{-1}$  by using Cayley-Hamilton theorem for the matrix **07**

$$A = \begin{bmatrix} 1 & 3 & 2 \\ 0 & -1 & 4 \\ -2 & 1 & 5 \end{bmatrix}. \text{ Hence find the matrix represented by}$$

**Q.7 (a)** Show that  $F = (y^2 - z^2 + 3yz - 2x)i + (3xz + 2xy)j + (3xy - 2xz + 2z)k$  is both solenoidal and irrotational. **03**

**(b)** Find the work done when a force  $F = (x^2 - y^2 + 2x)\hat{i} - (2xy + y)\hat{j}$  moves a particle in the  $xy$ -plane from  $(0,0)$  to  $(1,1)$  along the parabola  $y^2 = x$ . Is the work done different when the path is the straight line  $y = x$ ? **04**

**(c)** State Green's theorem and use it to evaluate the integral  $\oint_C y^2 dx + x^2 dy$ , where  $C$  is the triangle bounded by  $x = 0, x + y = 1, y = 0$ . **07**

\*\*\*\*\*